



Monthly Bulletin

Institut de physique du globe de Paris
Observatoire volcanologique du Piton de la Fournaise

ISSN 2610 – 5101

June, 2023

PITON DE LA FOURNAISE (VNUM #233020)

Latitude: 21.244°S

Longitude: 55.708°E

Summit elevation: 2632 m

Piton de la Fournaise is a basaltic hot spot volcano located in the southeastern part of La Réunion Island (Indian Ocean).

The volcano first erupted about 500,000 years ago. Its volcanic activity is characterized by frequent effusive eruptions (with emissions of lava fountains and lava flows) that occur on average twice a year since 1998. More rarely, larger explosive eruptions (with blocks covering the summit area and ash emissions that can disperse over long distances) have happened in the past with a centennial recurrence rate.

Most of the current eruptive activity (97% during the last 300 years) occurs from vents inside the Enclos Fouqué caldera. A few eruptions, however, have occurred from vents outside the caldera (most recently in 1977, 1986, and 1998). Such eruptions can potentially threaten communities that live in the surrounding areas.

Since late 1979, the activity of Piton de la Fournaise is monitored by the Piton de la Fournaise Volcanological Observatory (Observatoire Volcanologique du Piton de la Fournaise - OVPF), which belongs to the Institut de Physique du Globe de Paris (IPGP).

Alert level: Vigilance

(April 24, 2023 – July 2, 2023)

(cf. table in the appendix)



A. Piton de la Fournaise activity

Seismicity

In June 2023, the OVPF-IPGP recorded at Piton de La Fournaise:

- 791 shallow volcano-tectonic earthquakes (0 to 2.5 km above sea level) below the summit craters;
- 2 deep earthquakes (below sea level);
- 18 long-period earthquakes;
- 163 rockfalls.





Figure 1: Number of (top) shallow volcano-tectonic earthquakes and (bottom) rockfalls per day recorded in June 2023 (© OVPF-IPGP).

The seismic activity at Piton de la Fournaise in June was marked by an increase of the seismicity below the summit from June 12, from about 4 shallow volcano-tectonic earthquakes per day at the beginning of the month to 40 per day the last 15 days (Figure 1).

These earthquakes were located below the Dolomieu crater between -500 and 1200 m above sea level (Figure 2).

Numerous rockfalls also occurred inside the *Cratère Dolomieu*, along the cliffs of the *Rivière de l'Est* and on the September-October lava flows (Figure 1).




PROC.HYPO / Endos_ - sysop@pitondescalumets - 03-Jul-2023 19:10:08 +0 - hypomap.m (2021-05-19) / WebObs MMXXIII

Figure 2: Seismicity below Piton de la Fournaise in June 2023. Location map (epicenters) and north-south and east-west cross-sections (hypocenters) of earthquakes as recorded by OVPF-IPGP. Only manually located earthquakes are shown on the map (© OVPF-IPGP).




Deformation

Summit inflation, which had been stopped at the beginning of May, resumed in mid-June (Figures 3 and 4). The slow inflation (maximum elongation of the terminal cone < 2 cm in 15 days; Figure 4) is linked to the pressurization of the shallow magmatic reservoir located 1.5-2 km below the craters (Figure 5).



PROC.GIPSYX / BASELINES - sysop@pitondescalumets - 03-Jul-2023 19:01:44 +0 - gnss.m (2022-06-12) / WebObs M000XII

Figure 3: Ground deformation records over the past six months (in case of eruptive or intrusive periods, red and green bars represent eruptions and intrusions, respectively). The time series plots show the changes in distance between pairs of GPS stations located around the Dolomieu summit crater (reference: SNEG; top graph), the terminal cone (reference: FOAG; middle graph) and the Enclos Fouqué caldera (reference: PRAG; bottom graph), from north to south (see location in Figure 6). Increasing distances (or baseline elongation) indicate volcano inflation, while decreasing distances (or baseline contraction) reflect edifice deflation (© OVPF-IPGP).



PROC.GIPSYX / BASELINES_ - sysop@piton.descalumets - 03-Jul-2023 19:10:08 +0 - gnss.m (2022-06-12) / WebObs MMXXIII

Figure 4: Ground deformation records over the course of June 2023 (in case of eruptive or intrusive periods, red and green bars represent eruptions and intrusions, respectively). The time series plots show the changes in distance between pairs of GPS stations located around the Dolomieu summit crater (reference: SNEG; top graph), the terminal cone (reference: FOAG; middle graph) and the Enclos Fouqué caldera (reference: PRAG; bottom graph), from north to south (see location in Figure 6). Increasing distances (or baseline elongation) indicate volcano inflation, while decreasing distances (or baseline contraction) reflect edifice deflation (© OVPF-IPGP).



GNSS GIPSYX Pdf OVPF – Relative vectors
© OVPF-IPGP, 2023




Figure 5: Map of ground displacements (expressed in velocity) recorded in June 2023 by the OVPF-IPGP permanent GPS network. Horizontal displacements are represented in vector form and vertical displacements are indicated by numerical values in color (© OVPF-IPGP)




Figure 6: Location map of GPS stations and baselines as discussed in the text and shown in Figures 3 and 4 (© OVPF-IPGP).

* Glossary: The summit GPS signals indicate the influence of a shallow pressure source below the volcano, while distant GPS signals indicate the influence of a deep pressure source below the volcano. Inflation usually means pressurization; and conversely deflation usually means depressurization.



Gas geochemistry

CO₂ concentration in the soil

In the proximal Gite site, a sudden drop to very low CO₂ fluxes was detected after January 3, 2022. Since the end of the December 22 - January 17 eruption a new phase of increase was recorded, but with a lower rate.

The significant fluctuations observed during February 2022 are likely related to the environmental influence of two cyclonic events (Figure 7).




Figure 7: Comparison between the normalized average of uncorrected (OVPF-model; 15 days moving average; in green) and corrected for long-term influence of environmental parameters (INGV MALFIT model; in black) soil CO₂ flux from distal stations since October 2016 (last station set). Red bars: eruptions; Gray bars: intrusions (© OVPF-IPGP).

A new increase in soil CO₂ emissions was recorded in both distal and proximal stations at the end of February 2022 (Figure 7), with a strong acceleration from March 15. The new phase of increase in CO₂ soil emissions has lasted till May 05 in the distal area and till May 19, 2022 in the proximal area.

Since mid-May 2022, a trend of decrease in CO₂ gas fluxes is recorded in both proximal and distal sites. The September 19, 2022 eruption occurred after a significant decrease in CO₂ fluxes, likely recording the progressive transfer of magma to shallow crustal levels. Since the end of the September 19 – October 5 eruption CO₂ fluxes have remained on a stable level.

Interestingly, isotopic analysis of gas sampled at both distal (PNRN, BLEN, PCNR) and proximal (P0; GITN) sites shows a marked increase in the magmatic contribution in the March-April 2022 period (Figure 8). The magmatic contribution has then decreased in the second half of 2022.




Figure 8: Carbon isotope ($\delta^{13}C$) variations in CO_2 from soil gas emitted from the control point with the highest flux in the proximal area (control point: GIT0).

A new trend of increase in soil CO_2 emissions has started since the beginning of December 2022 in both distal and proximal sites. The increase has occurred with a high rate till mid-march 2023 and then it has slowed down.

A decrease in the CO_2 emission rate in the distal stations is measured since mid-June. And since June 23, 2023 the *Gîte* proximal station records a very strong decrease in CO_2 emission, heralding a possible transfer of magma to shallow depth.

* Glossary: CO_2 is the first gas to be released from deep magma (rising from the mantle), so its detection in the far field often means a deep rise of magma. Its near-field evolution may be related to magmatic transfer in the shallowest part of the feeding system (< 2-4 km below the surface).

Summit fumaroles composition obtained by the MultiGas method

- SO_2 : close or below the detection threshold.

* Glossary: The MultiGaS method allows measuring the concentrations of H_2O , H_2S , SO_2 and CO_2 in the atmosphere at the summit of the Piton de la Fournaise volcano. Magmatic transfer in the Piton de la Fournaise feeding system can result in an increase in SO_2 concentrations and in the C/S ratio (carbon/sulfur).



SO₂ flux in the air obtained by DOAS method

The SO₂ fluxes in the air are low; close or below the detection threshold.

** Glossary: During rest periods, SO₂ flux at Piton de la Fournaise is below the detection threshold. The SO₂ flux may increase during magma transfer in the shallowest part of the feeding system. During eruptions, it is directly proportional to the amount of lava emitted at the surface.*

Phenomenology

No eruptive activity reported in June 2023.

Summary

In June 2023, the pressurization of the shallow magma reservoir resumed from mid-June, with an inflation of the whole edifice and a mean of 40 shallow volcano-tectonic earthquakes per day. Since June 23, 2023 the *Gîte* proximal station records a very strong decrease in CO₂ emission, heralding a possible transfer of magma to shallow depth.



B. Seismic activity on La Réunion and in the Indian Ocean basin

Local and regional seismicity

In June 2023, the OVPF-IPGP recorded:

- 70 local earthquakes (below the island, within a radius of 200 km around the island, Figures 9 and 10);
- 1 regional earthquake (in the Indian Ocean basin).

In June, the OVPF-IPGP recorded 70 local earthquakes below the La Réunion island, and mainly near *Cirque de Salazie* (Figure 10).

These earthquakes were located between 10 km and 20 km depth in oceanic lithosphere on which was built the volcanic edifice at the origin of La Réunion.




Figure 9: Number of local earthquakes (La Réunion island) per day recorded in June 2023 (© OVPF-IPGP).




Figure 10: Seismicity below La Réunion in June 2023. Location map (epicenters) and north-west – south-east and south-west – north-east cross-sections (hypocenters) of earthquakes as recorded by OVPF-IPGP. Only localizable earthquakes are shown on the map (© OVPF-IPGP).



Seismic-volcano activity in Mayotte

The « REseau de surveillance VOlcanologique et Sismologique de MAyotte (REVOSIMA) » is the structure in charge of the volcano and seismic monitoring of Mayotte. IGP operates this network through the Piton de la Fournaise Volcanological Observatory in La Réunion with the support of the BRGM regional office in Mayotte. REVOSIMA is supported by a scientific and technical partnership. The REVOSIMA consortium: IGP and Université Paris Cité, BRGM, IFREMER, CNRS, BCSF-RéNaSS, ITES and Université de Strasbourg, IGN, ENS, SHOM, TAAF, Météo France, CNES, Université Grenoble Alpes and ISTerre, Université Clermont Auvergne, LMV and OPGC, Université de La Réunion, Université Paul Sabatier, Toulouse and GET-OMP, Université de la Rochelle, Université de Bretagne Occidentale, IRD and collaborators.

All information on the REVOSIMA and the activity in Mayotte can be found on the dedicated webpages:

- <http://www.ipgp.fr/fr/reseau-de-surveillance-volcanologique-sismologique-de-mayotte>
- <http://www.ipgp.fr/fr/actualites-reseau>
- <https://www.facebook.com/ReseauVolcanoSismoMayotte/>

July, 4 2023
OVPF-IPGP Director



C. Appendix

Definition of Volcanic Alert Levels for Piton de la Fournaise

from *disposition spécifique « Volcan Piton de la Fournaise » - arrêté n°2242*- Emergency plan set up by the department responsible for the protection of the population in the event of unrest or activity of the Piton de la Fournaise

• **“Vigilance”**: possible eruption in medium term (a few days or weeks) or presence of risks on the sector (rockfalls, increase of gas emissions, still hot lava flows...).

Access to the Enclos Fouqué caldera and to the summit volcano are allowed with restrictions.

• **“Alert 1”**: probable or imminent.

Access to the Enclos Fouqué caldera and to the summit are closed and prohibited.

• **“Alert 2”**: ongoing eruption.

Alert 2-1: ongoing eruption inside the Enclos Fouqué caldera without threat to the safety of people, property or the environment

Alert 2-2: ongoing eruption inside the Enclos Fouqué caldera with direct or indirect threat to the safety of people, property or the environment.

Access to the Enclos Fouqué caldera and to the summit are closed and prohibited. For Alert 2-2, evacuation of the people and vehicles depending on the issues.

• **“Alert 2-3”**: ongoing eruption outside the Enclos Fouqué caldera with threat to the safety of people, property or the environment.

Access to the Enclos Fouqué caldera and to the summit are closed and prohibited. Evacuation of the people and vehicles depending on the issues.

• **“Sauvegarde”**: end of eruption.

Evaluation of a partial reopening of the Enclos Fouqué caldera access.



Synthetic report on the last emitted products

To characterize the eruption of 19 September - 5 October 2022, 8 samples were collected among which two are pyroclast fall out deposits (lapilli, scoria) and the other 6 are lava fragments that were either collected in a molten state and quenched on site (syn-eruptive), or collected once solidified (post-emplacment).



Figure 11: Fragments of samples collected by the OVPF during the 19 September - 5 October 2022 eruption of Piton de la Fournaise

The pyroclastic fragments show a vesicularity of 37vol% to 76vol%, with up to 33vol% of isolated vesicles. The golden pumice scoria deposits are the most vesicular fragments (68-80 vol%) with significant values of isolated vesicles (up to 26vol%). The lava samples have a lower vesicularity (28-60vol%) than the pyroclasts. As previously observed, the quenched lavas have a lower vesicularity than the one collected post-emplacment and have some isolated vesicles (17vol%). **The data from this eruption confirm the increase in density of the eruptive products measured since 2019 and the overall trend of increasing density since the start of the eruptive cycle in 2014.**

All samples are nearly aphyric in terms of phenocrysts (<5vol% phenocrysts), with microphenocryst content ranging from 13vol% in pyroclasts to 21- 29vol.% in lavas.

The composition of the interstitial glassy matrix and microlites are similar to those of the lavas emitted during previous eruptions since 2020 and indicate a pre-eruptive temperature that varies between 1140°C at the beginning of the eruption to 1153°C towards the end of the eruption. **These relatively low temperatures are consistent with a shallow magmatic system.**

On long-term geochemical monitoring, the lavas emitted until the eruption of 22 December 2021 - 17 January 2022 have an intermediate CaO/Al₂O₃ signature - an indication of the degree of magma evolution - which decreases with the eruption of 19 September - 5 October 2022. **This trend could mark a progressive cooling of the shallow part of the magma system.**

As often observed during long-lasting eruptions, the lavas emitted at the end of the eruption (samples from 5 oct. 2022) are slightly more mafic (higher MgO, higher CaO/Al₂O₃ ratio and higher Ni and Cr content) and have slightly higher ratios of incompatible elements than those emitted at the beginning of the eruption. This temporal evolution is indicative of heterogeneities in the magma plumbing system. **These variations are correlated with a slight increase in the MgO concentration of the glasses, indicating a slightly hotter magma emitted towards the end of the eruption.**

The products of the September-October 2022 eruption have a 'depleted' signature (low La/Yb ratio), which has been attributed to the melting of a more refractory mantle source. **The decrease in the trace element signature of the eruption was attributed to the melting of a more refractory mantle source.**



Acknowledgments

Thank you to organizations, communities and associations for publicly posting this report for the widest dissemination

Information

All information on the Piton de la Fournaise activity can be found on the OVPF-IPGP media:

- Internet website : ipgp.fr/fr/ovpf/actualites-ovpf
- Twitter : twitter.com/obsfournaise
- Facebook : facebook.com/ObsVolcanoPitonFournaise

A preliminary automatic daily bulletin of the OVPF-IPGP, relating to the activities of the day before, validated by an analyst, is published daily. It can be accessed directly at this link:

http://volcano.ipgp.fr/reunion/Bulletin_quotidien/bulletin.html

The seismicity validated in continuous by OVPF-IPGP can also be followed on the RENASS portal: <https://renass.unis-tra.fr/fr/zones/la-reunion>

The information in this document may not be used without explicit reference.